

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68869 A3

- (51) International Patent Classification⁷: C12N 15/54, 9/12
- (21) International Application Number: PCT/US01/07500
- (22) International Filing Date: 8 March 2001 (08.03.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/188,449 10 March 2000 (10.03.2000) US
- (71) Applicant: LEXICON GENETICS INCORPORATED
[US/US]: 4000 Research Forest Drive, The Woodlands, TX 77381 (US).
- (72) Inventors: WALKE, D., Wade: 7507 Danehill Drive, Spring, TX 77389 (US). WILGANOWSKI, Nathaniel, L.: 505 Cypress Station #1906, Houston, TX 77090 (US). TURNER, C., Alexander, Jr.: 67 Winter Wheat Place, The Woodlands, TX 77381 (US).
- (74) Agents: ISHIMOTO, Lance, K. et al.; Lexicon Genetics Incorporated, 4000 Research Forest Drive, The Woodlands, TX 77381 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- (88) Date of publication of the international search report:
24 January 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/68869 A3

(54) Title: HUMAN G-COUPLED PROTEIN RECEPTOR KINSAES AND POLYNUCLEOTIDES ENCODING THE SAME

(57) Abstract: Novel human polynucleotide and polypeptide sequences are disclosed that can be used in therapeutic, diagnostic, and pharmacogenomic applications.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/07500

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/54 C12N9/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, SEQUENCE SEARCH, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 01 38503 A (PLOWMAN GREGORY D ;CLARY DOUGLAS (US); SUGEN INC (US); WHYTE DAVID) 31 May 2001 (2001-05-31) Sequence 2 figures 1B-C Sequence 59 figure 2A	1-4
P,X	--- DATABASE EMBL [Online] 7 May 2000 (2000-05-07) BIRREN B. ET AL.: "Homo sapiens chromosome 17, clone RP11-144C9" Database accession no. AC068693 XP002181811 nts.51942-52281, 86466-87092 --- -/--	1,2

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

1 November 2001

Date of mailing of the international search report

08. 11. 01

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Blanco Urgoiti, B

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/07500

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WEISS ELLEN R. ET AL: "The cloning of GRK7, a candidate cone opsin kinase, from cone and rod-dominant mammalian retinas." MOLECULAR VISION, vol. 4, 8 December 1998 (1998-12-08), page 27 XP002181808 ---	
A	PREMONT RICHARD T ET AL: "The GRK4 subfamily of G protein-coupled receptor kinases: Alternative splicing, gene organization, and sequence conservation." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 274, no. 41, 8 October 1999 (1999-10-08), pages 29381-29389, XP002181809 ISSN: 0021-9258 ---	
A	BENOVIC JEFFREY L ET AL: "Molecular cloning and expression of GRK6: A new member of the G protein-coupled receptor kinase family." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 268, no. 26, 1993, pages 19521-19527, XP002181810 ISSN: 0021-9258 -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/07500

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0138503 A	31-05-2001	AU WO 1926001 A 0138503 A2	04-06-2001 31-05-2001

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68869 A2

(51) International Patent Classification⁷: C12N 15/54, 9/12

(21) International Application Number: PCT/US01/07500

(22) International Filing Date: 8 March 2001 (08.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/188,449 10 March 2000 (10.03.2000) US

(71) Applicant: **LEXICON GENETICS INCORPORATED**
[US/US]; 4000 Research Forest Drive, The Woodlands, TX
77381 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(72) Inventors: **WALKE, D., Wade**; 7507 Danehill Drive, Spring, TX 77389 (US). **WILGANOWSKI, Nathaniel, L.**; 505 Cypress Station #1906, Houston, TX 77090 (US). **TURNER, C., Alexander, Jr.**; 67 Winter Wheat Place, The Woodlands, TX 77381 (US).

(74) Agents: **ISHIMOTO, Lance, K.** et al.; Lexicon Genetics Incorporated, 4000 Research Forest Drive, The Woodlands, TX 77381 (US).

(54) Title: NOVEL HUMAN G-COUPLED PROTEIN RECEPTOR KINASES AND POLYNUCLEOTIDES ENCODING THE SAME

(57) Abstract: Novel human polynucleotide and polypeptide sequences are disclosed that can be used in therapeutic, diagnostic, and pharmacogenomic applications.

WO 01/68869 A2

NOVEL HUMAN G-COUPLED PROTEIN RECEPTOR KINASES AND
POLYNUCLEOTIDES ENCODING THE SAME

The present application claims the benefit of U.S.
5 Provisional Application Number 60/188,449 which was filed on March
10, 2000 and is herein incorporated by reference in its entirety.

1. INTRODUCTION

The present invention relates to the discovery,
identification, and characterization of novel human
10 polynucleotides encoding proteins that share sequence similarity
with animal kinases. The invention encompasses the described
polynucleotides, host cell expression systems, the encoded
proteins, fusion proteins, polypeptides and peptides, antibodies
to the encoded proteins and peptides, and genetically engineered
15 animals that either lack or over express the disclosed
polynucleotides, antagonists and agonists of the proteins, and
other compounds that modulate the expression or activity of the
proteins encoded by the disclosed polynucleotides that can be used
for diagnosis, drug screening, clinical trial monitoring and the
20 treatment of diseases and physiological disorders.

2. BACKGROUND OF THE INVENTION

Kinases mediate phosphorylation of a wide variety of proteins
and compounds in the cell. In conjunction with phosphatases,
kinases are involved in a range of regulatory and signaling
25 pathways. Given the physiological importance of kinases, they
have been subject to intense scrutiny and are proven drug targets.

3. SUMMARY OF THE INVENTION

The present invention relates to the discovery,
identification, and characterization of nucleotides that encode
30 novel human proteins and the corresponding amino acid sequences of
these proteins. The novel human proteins (NHPs) described for the

first time herein share structural similarity with animal kinases, including, but not limited to G-protein coupled receptor kinases (GRKs). As such, the novel polynucleotides encode novel GRKs having homologues and orthologs across a range of phyla and species.

The novel human polynucleotides described herein, encode open reading frames (ORFs) encoding proteins of 553 and 353 amino acids in length (see SEQ ID NOS:2 and 4 respectively).

The invention also encompasses agonists and antagonists of the described NHPs, including small molecules, large molecules, mutant NHPs, or portions thereof, that compete with native NHP, peptides, and antibodies, as well as nucleotide sequences that can be used to inhibit the expression of the described NHPs (e.g., antisense and ribozyme molecules, and gene or regulatory sequence replacement constructs) or to enhance the expression of the described NHP polynucleotides (e.g., expression constructs that place the described polynucleotide under the control of a strong promoter system), and transgenic animals that express a NHP transgene, or "knock-outs" (which can be conditional) that do not express a functional NHP. Knock-out mice can be produced in several ways, one of which involves the use of mouse embryonic stem cells ("ES cells") lines that contain gene trap mutations in a murine homolog of at least one of the described NHPs. When the unique NHP sequences described in SEQ ID NOS:1-5 are "knocked-out" they provide a method of identifying phenotypic expression of the particular gene as well as a method of assigning function to previously unknown genes. Additionally, the unique NHP sequences described in SEQ ID NOS:1-5 are useful for the identification of coding sequence and the mapping a unique gene to a particular chromosome.

Further, the present invention also relates to processes for identifying compounds that modulate, i.e., act as agonists or antagonists, of NHP expression and/or NHP activity that utilize

purified preparations of the described NHPs and/or NHP product, or cells expressing the same. Such compounds can be used as therapeutic agents for the treatment of any of a wide variety of symptoms associated with biological disorders or imbalances.

5

4. DESCRIPTION OF THE SEQUENCE LISTING AND FIGURES

The Sequence Listing provides the sequence of the novel human ORFs encoding the described novel human kinase proteins. SEQ ID NO:5 describes a full length ORF and flanking regions.

10

5. DETAILED DESCRIPTION OF THE INVENTION

The NHPs, described for the first time herein, are novel proteins that are expressed in, *inter alia*, human cell lines, and human fetal brain, adult brain, pituitary, cerebellum, spinal
15 cord, thymus, kidney, fetal liver, prostate, testis, adrenal gland, small intestine, skeletal muscle, uterus, placenta, mammary gland, and pericardium cells. The described sequences were compiled from gene trapped sequences in conjunction with sequences available in GENBANK, and cDNAs from adrenal gland,
20 skeletal muscle, thymus, and testis libraries (Edge Biosystems, Gaithersburg, MD).

The present invention encompasses the nucleotides presented in the Sequence Listing, host cells expressing such nucleotides, the expression products of such nucleotides, and: (a) nucleotides
25 that encode mammalian homologs of the described polynucleotides, including the specifically described NHPs, and the NHP products; (b) nucleotides that encode one or more portions of an NHP that correspond to functional domains, and the polypeptide products specified by such nucleotide sequences, including but not limited
30 to the novel regions of any active domain(s); (c) isolated nucleotides that encode mutant versions, engineered or naturally occurring, of the described NHPs in which all or a part of at least one domain is deleted or altered, and the polypeptide

products specified by such nucleotide sequences, including but not limited to soluble proteins and peptides in which all or a portion of the signal sequence is deleted; (d) nucleotides that encode chimeric fusion proteins containing all or a portion of a coding region of a NHP, or one of its domains (e.g., a receptor/ligand binding domain, accessory protein/self-association domain, etc.) fused to another peptide or polypeptide; or (e) therapeutic or diagnostic derivatives of the described polynucleotides such as oligonucleotides, antisense polynucleotides, ribozymes, dsRNA, or gene therapy constructs comprising a sequence first disclosed in the Sequence Listing. As discussed above, the present invention includes: (a) the human DNA sequences presented in the Sequence Listing (and vectors comprising the same) and additionally contemplates any nucleotide sequence encoding a contiguous NHP open reading frame (ORF) that hybridizes to a complement of a DNA sequence presented in the Sequence Listing under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1xSSC/0.1% SDS at 68°C (Ausubel F.M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at p. 2.10.3) and encodes a functionally equivalent gene product. Additionally contemplated are any nucleotide sequences that hybridize to the complement of the DNA sequence that encode and express an amino acid sequence presented in the Sequence Listing under moderately stringent conditions, e.g., washing in 0.2xSSC/0.1% SDS at 42°C (Ausubel et al., 1989, *supra*), yet still encode a functionally equivalent NHP product. Functional equivalents of a NHP include naturally occurring NHPs present in other species and mutant NHPs whether naturally occurring or engineered (by site directed mutagenesis, gene shuffling, directed evolution as described in, for example, U.S. Patent No.

5,837,458). The invention also includes degenerate nucleic acid variants of the disclosed NHP polynucleotide sequences.

Additionally contemplated are polynucleotides encoding NHP ORFs, or their functional equivalents, encoded by polynucleotide
5 sequences that are about 99, 95, 90, or about 85 percent similar to corresponding regions of SEQ ID NO:1 (as measured by BLAST sequence comparison analysis using, for example, the GCG sequence analysis package using default parameters).

The invention also includes nucleic acid molecules,
10 preferably DNA molecules, that hybridize to, and are therefore the complements of, the described NHP encoding polynucleotides. Such hybridization conditions can be highly stringent or less highly stringent, as described above. In instances where the nucleic acid molecules are deoxyoligonucleotides ("DNA oligos"), such
15 molecules are generally about 16 to about 100 bases long, or about 20 to about 80, or about 34 to about 45 bases long, or any variation or combination of sizes represented therein that incorporate a contiguous region of sequence first disclosed in the Sequence Listing. Such oligonucleotides can be used in
20 conjunction with the polymerase chain reaction (PCR) to screen libraries, isolate clones, and prepare cloning and sequencing templates, etc..

Alternatively, such NHP oligonucleotides can be used as hybridization probes for screening libraries, and assessing gene
25 expression patterns (particularly using a micro array or high-throughput "chip" format). Additionally, a series of the described NHP oligonucleotide sequences, or the complements thereof, can be used to represent all or a portion of the described NHP sequences. An oligonucleotide or polynucleotide
30 sequence first disclosed in at least a portion of one or more of the sequences of SEQ ID NOS: 1-5 can be used as a hybridization probe in conjunction with a solid support matrix/substrate (resins, beads, membranes, plastics, polymers, metal or metallized

substrates, crystalline or polycrystalline substrates, etc.). Of particular note are spatially addressable arrays (*i.e.*, gene chips, microtiter plates, etc.) of oligonucleotides and polynucleotides, or corresponding oligopeptides and polypeptides, wherein at least one of the biopolymers present on the spatially addressable array comprises an oligonucleotide or polynucleotide sequence first disclosed in at least one of the sequences of SEQ ID NOS: 1-5, or an amino acid sequence encoded thereby. Methods for attaching biopolymers to, or synthesizing biopolymers on, solid support matrices, and conducting binding studies thereon are disclosed in, *inter alia*, U.S. Patent Nos. 5,700,637, 5,556,752, 5,744,305, 4,631,211, 5,445,934, 5,252,743, 4,713,326, 5,424,186, and 4,689,405 the disclosures of which are herein incorporated by reference in their entirety.

Addressable arrays comprising sequences first disclosed in SEQ ID NOS:1-5 can be used to identify and characterize the temporal and tissue specific expression of a gene. These addressable arrays incorporate oligonucleotide sequences of sufficient length to confer the required specificity, yet be within the limitations of the production technology. The length of these probes is within a range of between about 8 to about 2000 nucleotides. Preferably the probes consist of 60 nucleotides and more preferably 25 nucleotides from the sequences first disclosed in SEQ ID NOS:1-5.

For example, a series of the described oligonucleotide sequences, or the complements thereof, can be used in chip format to represent all or a portion of the described sequences. The oligonucleotides, typically between about 16 to about 40 (or any whole number within the stated range) nucleotides in length can partially overlap each other and/or the sequence may be represented using oligonucleotides that do not overlap. Accordingly, the described polynucleotide sequences shall typically comprise at least about two or three distinct

oligonucleotide sequences of at least about 8 nucleotides in length that are each first disclosed in the described Sequence Listing. Such oligonucleotide sequences can begin at any nucleotide present within a sequence in the Sequence Listing and
5 proceed in either a sense (5'-to-3') orientation vis-a-vis the described sequence or in an antisense orientation.

Microarray-based analysis allows the discovery of broad patterns of genetic activity, providing new understanding of gene functions and generating novel and unexpected insight into
10 transcriptional processes and biological mechanisms. The use of addressable arrays comprising sequences first disclosed in SEQ ID NOS:1-5 provides detailed information about transcriptional changes involved in a specific pathway, potentially leading to the identification of novel components or gene functions that manifest
15 themselves as novel phenotypes.

Probes consisting of sequences first disclosed in SEQ ID NOS:1-5 can also be used in the identification, selection and validation of novel molecular targets for drug discovery. The use of these unique sequences permits the direct confirmation of drug
20 targets and recognition of drug dependent changes in gene expression that are modulated through pathways distinct from the drugs intended target. These unique sequences therefore also have utility in defining and monitoring both drug action and toxicity.

As an example of utility, the sequences first disclosed in
25 SEQ ID NOS:1-5 can be utilized in microarrays or other assay formats, to screen collections of genetic material from patients who have a particular medical condition. These investigations can also be carried out using the sequences first disclosed in SEQ ID NOS:1-5 *in silico* and by comparing previously collected genetic
30 databases and the disclosed sequences using computer software known to those in the art.

Thus the sequences first disclosed in SEQ ID NOS:1-5 can be used to identify mutations associated with a particular disease and also as a diagnostic or prognostic assay.

Although the presently described sequences have been specifically described using nucleotide sequence, it should be appreciated that each of the sequences can uniquely be described using any of a wide variety of additional structural attributes, or combinations thereof. For example, a given sequence can be described by the net composition of the nucleotides present within a given region of the sequence in conjunction with the presence of one or more specific oligonucleotide sequence(s) first disclosed in the SEQ ID NOS: 1-5. Alternatively, a restriction map specifying the relative positions of restriction endonuclease digestion sites, or various palindromic or other specific oligonucleotide sequences can be used to structurally describe a given sequence. Such restriction maps, which are typically generated by widely available computer programs (e.g., the University of Wisconsin GCG sequence analysis package, SEQUENCHER 3.0, Gene Codes Corp., Ann Arbor, MI, etc.), can optionally be used in conjunction with one or more discrete nucleotide sequence(s) present in the sequence that can be described by the relative position of the sequence relative to one or more additional sequence(s) or one or more restriction sites present in the disclosed sequence.

For oligonucleotide probes, highly stringent conditions may refer, e.g., to washing in 6xSSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos). These nucleic acid molecules may encode or act as NHP gene antisense molecules, useful, for example, in NHP gene regulation (for and/or as antisense primers in amplification reactions of NHP gene nucleic acid sequences). With respect to NHP gene regulation, such techniques can be used to regulate biological functions. Further,

such sequences can be used as part of ribozyme and/or triple helix sequences that are also useful for NHP gene regulation, or as NHP regulating aptamers.

Inhibitory antisense or double stranded oligonucleotides can additionally comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 10 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 15 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 20 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide can also comprise at least one modified sugar moiety selected from the group including but not 25 limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide will comprise at least one modified phosphate backbone selected from the group consisting of a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a 30 methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide

forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al., 5 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue et al., 1987, FEBS Lett. 215:327-330). Alternatively, double stranded RNA can be used to disrupt the expression and function of a targeted NHP.

Oligonucleotides of the invention can be synthesized by 10 standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides can be synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), and methylphosphonate 15 oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:7448-7451), etc.

Low stringency conditions are well known to those of skill in the art, and will vary predictably depending on the specific 20 organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual (and periodic updates thereof), Cold Springs Harbor Press, N.Y.; and Ausubel et al., 1989, Current Protocols in Molecular Biology, 25 Green Publishing Associates and Wiley Interscience, N.Y.

Alternatively, suitably labeled NHP nucleotide probes can be used to screen a human genomic library using appropriately stringent conditions or by PCR. The identification and characterization of human genomic clones is helpful for 30 identifying polymorphisms (including, but not limited to, nucleotide repeats, microsatellite alleles, single nucleotide polymorphisms, or coding single nucleotide polymorphisms),

determining the genomic structure of a given locus/allele, and designing diagnostic tests. For example, sequences derived from regions adjacent to the intron/exon boundaries of the human gene can be used to design primers for use in amplification assays to
5 detect mutations within the exons, introns, splice sites (e.g., splice acceptor and/or donor sites), etc., that can be used in diagnostics and pharmacogenomics.

Further, a NHP gene homolog can be isolated from nucleic acid from an organism of interest by performing PCR using two
10 degenerate or "wobble" oligonucleotide primer pools designed on the basis of amino acid sequences within the NHP products disclosed herein. The template for the reaction may be total RNA, mRNA, and/or cDNA obtained by reverse transcription of mRNA prepared from, for example, human or non-human cell lines or
15 tissue known or suspected to express an allele of a NHP gene.

The PCR product can be subcloned and sequenced to ensure that the amplified sequences represent the sequence of the desired NHP gene. The PCR fragment can then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified
20 fragment can be labeled and used to screen a cDNA library, such as a bacteriophage cDNA library. Alternatively, the labeled fragment can be used to isolate genomic clones via the screening of a genomic library.

PCR technology can also be used to isolate full length cDNA
25 sequences. For example, RNA can be isolated, following standard procedures, from an appropriate cellular or tissue source (i.e., one known, or suspected, to express a NHP gene). A reverse transcription (RT) reaction can be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the
30 amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" using a standard terminal transferase reaction, the hybrid may be digested with RNase H, and second strand synthesis may then be primed with a

complementary primer. Thus, cDNA sequences upstream of the amplified fragment can be isolated. For a review of cloning strategies that can be used, see e.g., Sambrook et al., 1989, *supra*.

5 A cDNA encoding a mutant NHP gene can be isolated, for example, by using PCR. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known or suspected to be expressed in an individual putatively carrying a mutant NHP allele, and by
10 extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal gene. Using these two primers, the product is then amplified via PCR, optionally cloned into a suitable vector, and subjected to DNA
15 sequence analysis through methods well known to those of skill in the art. By comparing the DNA sequence of the mutant NHP allele to that of a corresponding normal NHP allele, the mutation(s) responsible for the loss or alteration of function of the mutant NHP gene product can be ascertained.

20 Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of or known to carry a mutant NHP allele (e.g., a person manifesting a NHP-associated phenotype such as, for example, immune disorders, obesity, high blood pressure, etc.), or a cDNA library can be constructed using
25 RNA from a tissue known, or suspected, to express a mutant NHP allele. A normal NHP gene, or any suitable fragment thereof, can then be labeled and used as a probe to identify the corresponding mutant NHP allele in such libraries. Clones containing mutant NHP gene sequences can then be purified and subjected to sequence
30 analysis according to methods well known to those skilled in the art.

 Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a

tissue known, or suspected, to express a mutant NHP allele in an individual suspected of or known to carry such a mutant allele. In this manner, gene products made by the putatively mutant tissue may be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against a normal NHP product, as described below. (For screening techniques, see, for example, Harlow, E. and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor, NY)

10 Additionally, screening can be accomplished by screening with labeled NHP fusion proteins, such as, for example, alkaline phosphatase-NHP or NHP-alkaline phosphatase fusion proteins. In cases where a NHP mutation results in an expressed gene product with altered function (e.g., as a result of a missense or a
15 frameshift mutation), polyclonal antibodies to a NHP are likely to cross-react with a corresponding mutant NHP gene product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well known in the art.

20 An additional application of the described novel human polynucleotide sequences is their use in the molecular mutagenesis/evolution of proteins that are at least partially encoded by the described novel sequences using, for example, polynucleotide shuffling or related methodologies. Such
25 approaches are described in U.S. Patents Nos. 5,830,721 and 5,837,458 which are herein incorporated by reference in their entirety.

 The invention also encompasses (a) DNA vectors that contain any of the foregoing NHP coding sequences and/or their complements
30 (i.e., antisense); (b) DNA expression vectors that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences (for example, baculo virus as described in U.S. Patent

No. 5,869,336 herein incorporated by reference); (c) genetically engineered host cells that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell; and (d) genetically engineered host cells that express an endogenous NHP gene under the control of an exogenously introduced regulatory element (*i.e.*, gene activation). As used herein, regulatory elements include, but are not limited to, inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include but are not limited to the cytomegalovirus (hCMV) immediate early gene, regulatable, viral elements (particularly retroviral LTR promoters), the early or late promoters of SV40 adenovirus, the *lac* system, the *trp* system, the *TAC* system, the *TRC* system, the major operator and promoter regions of phage lambda, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase (PGK), the promoters of acid phosphatase, and the promoters of the yeast α -mating factors.

Where, as in the present instance, some of the described NHP peptides or polypeptides are thought to be cytoplasmic proteins, expression systems can be engineered that produce soluble derivatives of a NHP (corresponding to a NHP extracellular and/or intracellular domains, or truncated polypeptides lacking one or more hydrophobic domains) and/or NHP fusion protein products (especially NHP-Ig fusion proteins, *i.e.*, fusions of a NHP domain to an IgFc), NHP antibodies, and anti-idiotypic antibodies (including Fab fragments) that can be used in therapeutic applications. Preferably, the above expression systems are engineered to allow the desired peptide or polypeptide to be recovered from the culture media.

The present invention also encompasses antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists and

agonists of a NHP, as well as compounds or nucleotide constructs that inhibit expression of a NHP gene (transcription factor inhibitors, antisense and ribozyme molecules, or gene or regulatory sequence replacement constructs), or promote the
5 expression of a NHP (e.g., expression constructs in which NHP coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, etc.).

The NHPs or NHP peptides, NHP fusion proteins, NHP nucleotide sequences, antibodies, antagonists and agonists can be useful for
10 the detection of mutant NHPs or inappropriately expressed NHPs for the diagnosis of disease. The NHP proteins or peptides, NHP fusion proteins, NHP nucleotide sequences, host cell expression systems, antibodies, antagonists, agonists and genetically engineered cells and animals can be used for screening for drugs
15 (or high throughput screening of combinatorial libraries) effective in the treatment of the symptomatic or phenotypic manifestations of perturbing the normal function of a NHP in the body. The use of engineered host cells and/or animals can offer an advantage in that such systems allow not only for the
20 identification of compounds that bind to the endogenous receptor/ligand of a NHP, but can also identify compounds that trigger NHP-mediated activities or pathways.

Finally, the NHP products can be used as therapeutics. For example, soluble derivatives such as NHP peptides/domains
25 corresponding to NHPs, NHP fusion protein products (especially NHP-Ig fusion proteins, i.e., fusions of a NHP, or a domain of a NHP, to an IgFc), NHP antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists (including compounds that modulate or act on downstream targets in a NHP-
30 mediated pathway) can be used to directly treat diseases or disorders. For instance, the administration of an effective amount of soluble NHP, or a NHP-IgFc fusion protein or an anti-idiotypic antibody (or its Fab) that mimics the NHP could activate

or effectively antagonize the endogenous NHP or a protein interactive therewith. Nucleotide constructs encoding such NHP products can be used to genetically engineer host cells to express such products *in vivo*; these genetically engineered cells function as "bioreactors" in the body delivering a continuous supply of a NHP, a NHP peptide, or a NHP fusion protein to the body.

Nucleotide constructs encoding functional NHPs, mutant NHPs, as well as antisense and ribozyme molecules can also be used in "gene therapy" approaches for the modulation of NHP expression. Thus, the invention also encompasses pharmaceutical formulations and methods for treating biological disorders.

Various aspects of the invention are described in greater detail in the subsections below.

5.1 THE NHP SEQUENCES

The cDNA sequences and corresponding deduced amino acid sequences of the described NHPs are presented in the Sequence Listing.

Expression analysis has provided evidence that the described NHPs can be expressed in human tissues as well as gene trapped human cells. In addition to GRKs, the described NHPs share significant similarity to a range of kinase families from a variety of phyla and species. Similar polynucleotides encoding GRK proteins, as well as uses and applications that are germane to the described NHPs are described in U.S. Patents Nos. 5,591,618 and 5,532,151 which are herein incorporated by reference in their entirety.

5.2 NHPS AND NHP POLYPEPTIDES

NHP products, polypeptides, peptide fragments, mutated, truncated, or deleted forms of the NHPs, and/or NHP fusion proteins can be prepared for a variety of uses. These uses include, but are not limited to, the generation of antibodies, as

reagents in diagnostic assays, the identification of other cellular gene products related to the NHP, as reagents in assays for screening for compounds that can be used as pharmaceutical reagents useful in the therapeutic treatment of mental,

5 biological, or medical disorders and disease.

The Sequence Listing discloses the amino acid sequences encoded by the described NHP-encoding polynucleotides. The NHPs display initiator methionines in a DNA sequence context consistent with eucaryotic translation initiation site, and a weak signal
10 sequence characteristic of membrane associated proteins.

The NHP amino acid sequences of the invention include the amino acid sequences presented in the Sequence Listing as well as analogues and derivatives thereof. Further, corresponding NHP homologues from other species are encompassed by the invention.
15 In fact, any NHP protein encoded by the NHP nucleotide sequences described above are within the scope of the invention, as are any novel polynucleotide sequences encoding all or any novel portion of an amino acid sequence presented in the Sequence Listing. The degenerate nature of the genetic code is well known, and,
20 accordingly, each amino acid presented in the Sequence Listing, is generically representative of the well known nucleic acid "triplet" codon, or in many cases codons, that can encode the amino acid. As such, as contemplated herein, the amino acid sequences presented in the Sequence Listing, when taken together
25 with the genetic code (see, for example, Table 4-1 at page 109 of "Molecular Cell Biology", 1986, J. Darnell et al. eds., Scientific American Books, New York, NY, herein incorporated by reference) are generically representative of all the various permutations and combinations of nucleic acid sequences that can encode such amino
30 acid sequences.

The invention also encompasses proteins that are functionally equivalent to the NHPs encoded by the presently described nucleotide sequences as judged by any of a number of criteria,

including, but not limited to, the ability to bind and modify a NHP substrate, or the ability to effect an identical or complementary downstream pathway, or a change in cellular metabolism (e.g., proteolytic activity, ion flux, tyrosine phosphorylation, etc.). Such functionally equivalent NHP proteins include, but are not limited to, additions or substitutions of amino acid residues within the amino acid sequence encoded by a NHP nucleotide sequence described above, but which result in a silent change, thus producing a functionally equivalent gene product. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

A variety of host-expression vector systems can be used to express the NHP nucleotide sequences of the invention. Where the NHP peptide or polypeptide can exist, or has been engineered to exist, as a soluble or secreted molecule, the soluble NHP peptide or polypeptide can be recovered from the culture media. Such expression systems also encompass engineered host cells that express a NHP, or functional equivalent, *in situ*. Purification or enrichment of a NHP from such expression systems can be accomplished using appropriate detergents and lipid micelles and methods well known to those skilled in the art. However, such engineered host cells themselves may be used in situations where it is important not only to retain the structural and functional characteristics of the NHP, but to assess biological activity, e.g., in drug screening assays.

The expression systems that may be used for purposes of the invention include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors
5 containing NHP nucleotide sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing NHP nucleotide sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing NHP sequences; plant cell systems infected with
10 recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing NHP nucleotide sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression
15 constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

In bacterial systems, a number of expression vectors may be
20 advantageously selected depending upon the use intended for the NHP product being expressed. For example, when a large quantity of such a protein is to be produced for the generation of pharmaceutical compositions of or containing NHP, or for raising antibodies to a NHP, vectors that direct the expression of high
25 levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which a NHP coding sequence may be ligated individually into the vector in frame with the *lacZ* coding region
30 so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke & Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX

vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by
5 elution in the presence of free glutathione. The PGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

In an insect system, *Autographa californica* nuclear
10 polyhidrosis virus (AcNPV) is used as a vector to express foreign polynucleotides. The virus grows in *Spodoptera frugiperda* cells. A NHP encoding polynucleotide sequence can be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for
15 example the polyhedrin promoter). Successful insertion of NHP coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus, lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera*
20 *frugiperda* cells in which the inserted polynucleotide is expressed (e.g., see Smith et al., 1983, J. Virol. 46: 584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as
25 an expression vector, the NHP nucleotide sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene can then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-
30 essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing a NHP product in infected hosts (e.g., See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific

initiation signals may also be required for efficient translation of inserted NHP nucleotide sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire NHP gene or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of a NHP coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (See Bitter *et al.*, 1987, *Methods in Enzymol.* 153:516-544).

In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in particular, human cell lines.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines that stably express the NHP sequences described above can be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the NHP product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the NHP product.

A number of selection systems can be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, Cell 22:817) genes can be employed in tk⁻, hgp⁻ or apt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA 77:3567; O'Hare, et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al.,

1981, J. Mol. Biol. 150:1); and hygromycin, which confers resistance to hygromycin (Santerre, et al., 1984, Gene 30:147).

Alternatively, any fusion protein can be readily purified by utilizing an antibody specific for the fusion protein being expressed. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl. Acad. Sci. USA 88:8972-8976). In this system, the polynucleotide of interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni²⁺ nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

Also encompassed by the present invention are fusion proteins that direct the NHP to a target organ and/or facilitate transport across the membrane into the cytosol. Conjugation of NHPs to antibody molecules or their Fab fragments could be used to target cells bearing a particular epitope. Attaching the appropriate signal sequence to the NHP would also transport the NHP to the desired location within the cell. Alternatively targeting of NHP or its nucleic acid sequence might be achieved using liposome or lipid complex based delivery systems. Such technologies are described in Liposomes: A Practical Approach, New, RRC ed., Oxford University Press, New York and in U.S. Patents Nos. 4,594,595, 5,459,127, 5,948,767 and 6,110,490 and their respective disclosures which are herein incorporated by reference in their entirety. Additionally embodied are novel protein constructs engineered in such a way that they facilitate transport of the NHP to the target site or desired organ, where they cross the cell membrane and/or the nucleus where the NHP can exert its functional activity. This goal may be achieved by coupling of the NHP to a

cytokine or other ligand that provides targeting specificity, and/or to a protein transducing domain (see generally U.S. applications Ser. No. 60/111,701 and 60/056,713, both of which are herein incorporated by reference, for examples of such transducing sequences) to facilitate passage across cellular membranes and can optionally be engineered to include nuclear localization sequences.

5.3 ANTIBODIES TO NHP PRODUCTS

10 Antibodies that specifically recognize one or more epitopes of a NHP, or epitopes of conserved variants of a NHP, or peptide fragments of a NHP are also encompassed by the invention. Such antibodies include but are not limited to polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, 15 single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

 The antibodies of the invention can be used, for example, in 20 the detection of NHP in a biological sample and may, therefore, be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal amounts of NHP. Such antibodies may also be utilized in conjunction with, for example, compound screening schemes for the evaluation of the effect of 25 test compounds on expression and/or activity of a NHP gene product. Additionally, such antibodies can be used in conjunction gene therapy to, for example, evaluate the normal and/or engineered NHP-expressing cells prior to their introduction into the patient. Such antibodies may additionally be used as a method 30 for the inhibition of abnormal NHP activity. Thus, such antibodies may, therefore, be utilized as part of treatment methods.

For the production of antibodies, various host animals may be immunized by injection with the NHP, a NHP peptide (e.g., one corresponding to a functional domain of a NHP), truncated NHP polypeptides (NHP in which one or more domains have been deleted), functional equivalents of the NHP or mutated variant of the NHP. Such host animals may include but are not limited to pigs, rabbits, mice, goats, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's adjuvant (complete and incomplete), mineral salts such as aluminum hydroxide or aluminum phosphate, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Alternatively, the immune response could be enhanced by combination and or coupling with molecules such as keyhole limpet hemocyanin, tetanus toxoid, diphtheria toxoid, ovalbumin, cholera toxin or fragments thereof. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of the immunized animals.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, can be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein, (1975, Nature 256:495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal Antibodies And Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated *in vitro* or

in vivo. Production of high titers of mAbs *in vivo* makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison *et al.*, 1984, Proc. Natl. Acad. Sci., 81:6851-6855; Neuberger *et al.*, 1984, Nature, 312:604-608; Takeda *et al.*, 1985, Nature, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region. Such technologies are described in U.S. Patents Nos. 6,075,181 and 5,877,397 and their respective disclosures which are herein incorporated by reference in their entirety. Also encompassed by the present invention is the use of fully humanized monoclonal antibodies as described in US Patent No. 6,150,584 and respective disclosures which are herein incorporated by reference in their entirety.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778; Bird, 1988, Science 242:423-426; Huston *et al.*, 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward *et al.*, 1989, Nature 341:544-546) can be adapted to produce single chain antibodies against NHP gene products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, such fragments include, but are not limited to: the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression

libraries may be constructed (Huse *et al.*, 1989, *Science*, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Antibodies to a NHP can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" a given NHP, using techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, 1993, *FASEB J* 7(5):437-444; and Nissinoff, 1991, *J. Immunol.* 147(8):2429-2438). For example antibodies which bind to a NHP domain and competitively inhibit the binding of NHP to its cognate receptor/ligand can be used to generate anti-idiotypes that "mimic" the NHP and, therefore, bind, activate, or neutralize a NHP, NHP receptor, or NHP ligand. Such anti-idiotypic antibodies or Fab fragments of such anti-idiotypes can be used in therapeutic regimens involving a NHP mediated pathway.

The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims. All cited publications, patents, and patent applications are herein incorporated by reference in their entirety.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising at least 24 contiguous bases of nucleotide sequence first disclosed in SEQ ID NO: 1.
2. An isolated nucleic acid molecule comprising a nucleotide sequence that:
 - (a) encodes the amino acid sequence shown in SEQ ID NO: 2 ; and
 - (b) hybridizes under stringent conditions to the nucleotide sequence of SEQ ID NO: 1 or the complement thereof.
3. An isolated nucleic acid molecule comprising a nucleotide sequence encoding the amino acid sequence shown in SEQ ID NO:2.
4. An isolated nucleic acid molecule comprising a nucleotide sequence encoding the amino acid sequence shown in SEQ ID NO:4.

SEQUENCE LISTING

<110> LEXICON GENETICS INCORPORATED

<120> Novel Human G-Coupled Protein Receptor Kinases and Polynucleotides
Encoding the Same

<130> LEX-0147-PCT

<150> US 60/188,449

<151> 2000-03-10

<160> 5

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1662

<212> DNA

<213> homo sapiens

<400> 1

atggtggaca	tgggggccct	ggayaacctg	atcgccaaca	ccgcctacct	gcaggcccg	60
aagccctcgg	actgcgacag	caaagagctg	cagcggcggc	ggcgtagcct	ggccctgccc	120
gggctgcagg	gctgcgcgga	gctccgccag	aagctgtccc	tgaacttcca	cagcctgtgt	180
gagcagcagc	ccatcggtcg	ccgcctcttc	cgtgacttcc	tagccacagt	gcccacgttc	240
cgcaaggcgg	caaccttcct	agaggacgtg	cagaactggg	agctggccga	ggagggaccc	300
accaaagaca	gcgcgctgca	ggggctggtg	gccacttgtg	cgagtgtccc	tgccccgggg	360
aaccgcgaac	ccttcctcag	ccaggccgtg	gccaccaagt	gccaagcagc	caccactgag	420
gaagagcgag	tggctgcagt	gacgctggcc	aaggctgagg	ccatggcttt	cttgcaagag	480
cagcccttta	aggatttcgt	gaccagcgcc	ttctacgaca	agtttctgca	gtggaaactc	540
ttcgagatgc	aaccagtgtc	agacaagtac	ttcactgagt	tcagagtgtc	ggggaaaggt	600
ggttttgagg	aggtatgtgc	cgtccaggtg	aaaaacactg	ggaagatgta	tgcctgtaa	660
aaactggaca	agaagcggct	gaagaagaaa	ggtggcgaga	agatggctct	cttggaanaa	720
gaaatcttgg	agaaggctcag	cagccctttc	attgtctctc	tggcctatgc	ctttgagagc	780
aagacccatc	tctgccttgt	catgagcctg	atgaatgggg	gagacctcaa	gttccacatc	840
tacaacgtgg	gcacgcgtgg	cctggacatg	agccgggtga	tcttttactc	ggcccagata	900
gcctgtggga	tgtctcacct	ccatgaactc	ggcatcgtct	atcgggacat	gaagcctgag	960
aatgtgtctc	tggatgacct	cggcaactgc	aggttatctg	acctggggct	ggccgtggag	1020
atgaagggtg	gcaagcccat	caccagagg	gctggaacca	atggttacat	ggctcctgag	1080
atcctaattg	aaaaggtaag	ttattcctat	cctgtggact	ggtttgccat	gggatgcagc	1140
atztatgaaa	tggttgctgg	acgaacacca	ttcaaagatt	acaaggaaaa	ggtcagtaaa	1200
gaggatctga	agcaaagaac	tctgcaagac	gaggtcataa	tccagcatga	taacttcaca	1260
gaggaagcaa	aagatatttg	caggctcttc	ttggctaaga	aaccagagca	acgcttagga	1320
agcagagaaa	agtctgatga	tcccaggaaa	catcatttct	ttaaaacgat	caactttcct	1380
cgcttggaag	ctggcctaata	tgaaccccca	tttgtgccag	acccttcagt	ggtttatgcc	1440
aaagacatcg	ctgaaattga	tgatttctct	gaggttcggg	gggtggaatt	tgatgacaaa	1500
gataagcagt	tcttcaaaaa	ctttgcgaca	ggtgtgttgc	ctatagcatg	gcaggaagaa	1560
attatagaaa	cgggactgtt	tgaggaactg	aatgacccca	acagacctac	gggttgtgag	1620
gagggttaatt	catccaagtc	tggcgtgtgt	ttgttattgt	aa		1662

<210> 2

<211> 553

<212> PRT

<213> homo sapiens

<400> 2

```

Met Val Asp Met Gly Ala Leu Asp Asn Leu Ile Ala Asn Thr Ala Tyr
 1           5           10          15
Leu Gln Ala Arg Lys Pro Ser Asp Cys Asp Ser Lys Glu Leu Gln Arg
 20          25          30
Arg Arg Arg Ser Leu Ala Leu Pro Gly Leu Gln Gly Cys Ala Glu Leu
 35          40          45
Arg Gln Lys Leu Ser Leu Asn Phe His Ser Leu Cys Glu Gln Gln Pro
 50          55          60
Ile Gly Arg Arg Leu Phe Arg Asp Phe Leu Ala Thr Val Pro Thr Phe
 65          70          75          80
Arg Lys Ala Ala Thr Phe Leu Glu Asp Val Gln Asn Trp Glu Leu Ala
 85          90          95
Glu Glu Gly Pro Thr Lys Asp Ser Ala Leu Gln Gly Leu Val Ala Thr
100         105         110
Cys Ala Ser Ala Pro Ala Pro Gly Asn Pro Gln Pro Phe Leu Ser Gln
115         120         125
Ala Val Ala Thr Lys Cys Gln Ala Ala Thr Thr Glu Glu Glu Arg Val
130         135         140
Ala Ala Val Thr Leu Ala Lys Ala Glu Ala Met Ala Phe Leu Gln Glu
145         150         155         160
Gln Pro Phe Lys Asp Phe Val Thr Ser Ala Phe Tyr Asp Lys Phe Leu
165         170         175
Gln Trp Lys Leu Phe Glu Met Gln Pro Val Ser Asp Lys Tyr Phe Thr
180         185         190
Glu Phe Arg Val Leu Gly Lys Gly Phe Gly Glu Val Cys Ala Val
195         200         205
Gln Val Lys Asn Thr Gly Lys Met Tyr Ala Cys Lys Lys Leu Asp Lys
210         215         220
Lys Arg Leu Lys Lys Lys Gly Gly Glu Lys Met Ala Leu Leu Glu Lys
225         230         235         240
Glu Ile Leu Glu Lys Val Ser Ser Pro Phe Ile Val Ser Leu Ala Tyr
245         250         255
Ala Phe Glu Ser Lys Thr His Leu Cys Leu Val Met Ser Leu Met Asn
260         265         270
Gly Gly Asp Leu Lys Phe His Ile Tyr Asn Val Gly Thr Arg Gly Leu
275         280         285
Asp Met Ser Arg Val Ile Phe Tyr Ser Ala Gln Ile Ala Cys Gly Met
290         295         300
Leu His Leu His Glu Leu Gly Ile Val Tyr Arg Asp Met Lys Pro Glu
305         310         315         320
Asn Val Leu Leu Asp Asp Leu Gly Asn Cys Arg Leu Ser Asp Leu Gly
325         330         335
Leu Ala Val Glu Met Lys Gly Gly Lys Pro Ile Thr Gln Arg Ala Gly
340         345         350
Thr Asn Gly Tyr Met Ala Pro Glu Ile Leu Met Glu Lys Val Ser Tyr
355         360         365
Ser Tyr Pro Val Asp Trp Phe Ala Met Gly Cys Ser Ile Tyr Glu Met
370         375         380
Val Ala Gly Arg Thr Pro Phe Lys Asp Tyr Lys Glu Lys Val Ser Lys
385         390         395         400
Glu Asp Leu Lys Gln Arg Thr Leu Gln Asp Glu Val Lys Phe Gln His
405         410         415
Asp Asn Phe Thr Glu Glu Ala Lys Asp Ile Cys Arg Leu Phe Leu Ala

```

420 425 430
 Lys Lys Pro Glu Gln Arg Leu Gly Ser Arg Glu Lys Ser Asp Asp Pro
 435 440 445
 Arg Lys His His Phe Phe Lys Thr Ile Asn Phe Pro Arg Leu Glu Ala
 450 455 460
 Gly Leu Ile Glu Pro Pro Phe Val Pro Asp Pro Ser Val Val Tyr Ala
 465 470 475 480
 Lys Asp Ile Ala Glu Ile Asp Asp Phe Ser Glu Val Arg Gly Val Glu
 485 490 495
 Phe Asp Asp Lys Asp Lys Gln Phe Phe Lys Asn Phe Ala Thr Gly Ala
 500 505 510
 Val Pro Ile Ala Trp Gln Glu Glu Ile Ile Glu Thr Gly Leu Phe Glu
 515 520 525
 Glu Leu Asn Asp Pro Asn Arg Pro Thr Gly Cys Glu Glu Gly Asn Ser
 530 535 540
 Ser Lys Ser Gly Val Cys Leu Leu Leu
 545 550

<210> 3

<211> 1062

<212> DNA

<213> homo sapiens

<400> 3

atggtggaca	tgggggccct	ggacaacctg	atcgccaaca	ccgcctacct	gcaggccccg	60
aagccctcgg	actgcgacag	caaagagctg	cagcggcggc	ggcgtagcct	ggccctgccc	120
gggtgcagg	gctgcgcgga	gctccgccag	aagctgtccc	tgaacttcca	cagcctgtgt	180
gagcagcagc	ccatcggtcg	ccgcctcttc	cgtgacttcc	tagccacagt	gccacggtc	240
cgcaaggcgg	caaccttctc	agaggacgtg	cagaactggg	agctggccga	ggagggacct	300
accaaagaca	gcgcgctgca	ggggctgggt	gccacttgtg	cgagtgcgcc	tgccccgggg	360
aaccgcgaac	ccttcctcag	ccaggccgtg	gccaccaagt	gccaagcagc	caccactgag	420
gaagagcgag	tggctgcagt	gacgctggcc	aaggctgagg	ccatggcttt	cttgcaagag	480
cagcccttta	aggatttcgt	gaccagcgcc	ttctacgaca	agtttctgca	gtggaaactc	540
ttcgagatgc	aaccagtgtc	agacaagtac	ttcactgagt	tcagagtgtc	ggggaaaggt	600
ggttttgagg	aggtatgtgc	cgtccagggt	aaaaaactg	ggaagatgta	tgctgtgaag	660
aaactggaca	agaagcggct	gaagaagaaa	ggtggcgaga	agatggctct	cttgaaaaag	720
gaaatcttgg	agaaggtcag	cagccctttc	attgtctctc	tggcctatgc	ctttgagagc	780
aagaccatc	tctgccttgt	catgagcctg	atgaatgggg	gagacctcaa	gttccacatc	840
tacaacgtgg	gcacgcgtgg	cctggacatg	agccgggtga	tcttttactc	ggcccagata	900
gcctgtggga	tgctgcacct	ccatgaactc	ggcatcgctc	atcgggacat	gaagcctgag	960
aatgtgcttc	tgatgacact	cggcaactgc	aggttatctg	acctggggct	ggccgtggag	1020
atgaagggtg	gcaagcccat	caccagagg	agaaaagtct	ga		1062

<210> 4

<211> 353

<212> PRT

<213> homo sapiens

<400> 4

Met Val Asp Met Gly Ala Leu Asp Asn Leu Ile Ala Asn Thr Ala Tyr
 1 5 10 15
 Leu Gln Ala Arg Lys Pro Ser Asp Cys Asp Ser Lys Glu Leu Gln Arg
 20 25 30
 Arg Arg Arg Ser Leu Ala Leu Pro Gly Leu Gln Gly Cys Ala Glu Leu
 35 40 45
 Arg Gln Lys Leu Ser Leu Asn Phe His Ser Leu Cys Glu Gln Gln Pro

50	55	60
Ile Gly Arg Arg Leu Phe Arg Asp Phe Leu Ala Thr Val Pro Thr Phe		
65	70	75
Arg Lys Ala Ala Thr Phe Leu Glu Asp Val Gln Asn Trp Glu Leu Ala		80
	85	90
Glu Glu Gly Pro Thr Lys Asp Ser Ala Leu Gln Gly Leu Val Ala Thr		95
	100	105
Cys Ala Ser Ala Pro Ala Pro Gly Asn Pro Gln Pro Phe Leu Ser Gln		110
	115	120
Ala Val Ala Thr Lys Cys Gln Ala Ala Thr Thr Glu Glu Glu Arg Val		125
	130	135
Ala Ala Val Thr Leu Ala Lys Ala Glu Ala Met Ala Phe Leu Gln Glu		140
145	150	155
Gln Pro Phe Lys Asp Phe Val Thr Ser Ala Phe Tyr Asp Lys Phe Leu		160
	165	170
Gln Trp Lys Leu Phe Glu Met Gln Pro Val Ser Asp Lys Tyr Phe Thr		175
	180	185
Glu Phe Arg Val Leu Gly Lys Gly Gly Phe Gly Glu Val Cys Ala Val		190
	195	200
Gln Val Lys Asn Thr Gly Lys Met Tyr Ala Cys Lys Lys Leu Asp Lys		205
	210	215
Lys Arg Leu Lys Lys Lys Gly Gly Glu Lys Met Ala Leu Leu Glu Lys		220
225	230	235
Glu Ile Leu Glu Lys Val Ser Ser Pro Phe Ile Val Ser Leu Ala Tyr		240
	245	250
Ala Phe Glu Ser Lys Thr His Leu Cys Leu Val Met Ser Leu Met Asn		255
	260	265
Gly Gly Asp Leu Lys Phe His Ile Tyr Asn Val Gly Thr Arg Gly Leu		270
	275	280
Asp Met Ser Arg Val Ile Phe Tyr Ser Ala Gln Ile Ala Cys Gly Met		285
	290	295
Leu His Leu His Glu Leu Gly Ile Val Tyr Arg Asp Met Lys Pro Glu		300
305	310	315
Asn Val Leu Leu Asp Asp Leu Gly Asn Cys Arg Leu Ser Asp Leu Gly		320
	325	330
Leu Ala Val Glu Met Lys Gly Gly Lys Pro Ile Thr Gln Arg Arg Lys		335
	340	345
Val		350

<210> 5

<211> 2249

<212> DNA

<213> homo sapiens

<400> 5

aaaactgctc tgaggccatc atgctttgag gaagcccagg aggaaacact gcagagaggc	60
tcaaccaccc cagctctccc agctgagctc agccaccac cgatcccca gctgaatgca	120
accataagag tgagtccagg ttctaccctg ctaggctgcc accacattcc taagaaccac	180
gggaaaaggc atttgctcct ccgaagaaat tctcagactg atttttcact gtattgtcag	240
gccacaggac tcaactgtaaa tcccttgac gttgtctcac ccgggaaggg aaagcagcca	300
gcagccctcc agccctcttg tgctttccct gggagtgcgc cccgtgctca gccatggtgg	360
acatgggggc cctggacaac ctgatcgcca acaccgccta cctgcaggcc cggaagccct	420
cggactgcga cagcaaagag ctgcagcggc ggcggcgtag cctggccctg cccgggctgc	480
agggctgcgc ggagctccgc cagaagctgt ccctgaactt ccacagcctg tgtgagcagc	540
agcccatcgg tcgccgcctc ttccgtgact tcctagccac agtgcccacg ttccgcaagg	600

cggaacac	cctagagg	gtgcagaact	gggagctggc	cgaggaggga	cccaccaaag	660
acagcgcgct	gcaggggctg	gtggccactt	gtgcgagtgc	ccctgccccg	gggaacccgc	720
aacccttcct	cagccaggcc	gtggccacca	agtgccaaagc	agccaccact	gaggaagagc	780
gagtggctgc	agtgcgctg	gccaaggctg	aggccatggc	tttcttgcaa	gagcagccct	840
ttaaggattt	cgtgaccagc	gccttctacg	acaagtttct	gcagtggaaa	ctcttcgaga	900
tgcaaccagt	gtcagacaag	tacttcaactg	agttcagagt	gctggggaaa	ggtgggtttg	960
gggaggtatg	tgccgtccag	gtgaaaaaca	ctgggaagat	gtatgcctgt	aagaaactgg	1020
acaagaagcg	gctgaagaag	aaaggtggcg	agaagatggc	tctcttgga	aaggaaatct	1080
tggaagaggt	cagcagccct	ttcattgtct	ctctggccta	tgcccttgag	agcaagaccc	1140
atctctgcct	tgatcatgagc	ctgatgaatg	ggggagacct	caagttccac	atctacaacg	1200
tgggcacgcg	tgccctggac	atgagccggg	tgatctttta	ctcgccccag	atagcctgtg	1260
ggatgctgca	cctccatgaa	ctcgcatcg	tctatcgga	catgaagcct	gagaatgtgc	1320
ttctggatga	cctcggaac	tgaggttat	ctgacctggg	gctggccgtg	gagatgaagg	1380
gtggcaagcc	catcacccag	agggtggaa	ccaatggtta	catggctcct	gagatcctaa	1440
tggaagaggt	aagttattcc	tatcctgtgg	actggtttgc	catgggatgc	agcatttatg	1500
aaatggttgc	tggaagcaaa	ccattcaaag	attacaagga	aaaggtcagt	aaagaggatc	1560
tgaagcaaa	aactctgcaa	gacgaggtca	aattccagca	tgataacttc	acagaggaag	1620
caaaagatat	ttgcaggctc	ttcttggcta	agaaaccaga	gcaacgctta	ggaagcagag	1680
aaaagtctga	tgatcccagg	aaacatcatt	tctttaaacc	gatcaacttt	cctcgccctg	1740
aagctggcct	aattgaaccc	ccatttgtgc	cagacccttc	agtggtttat	gcaaagaca	1800
tcgctgaaat	tgatgatattc	tctgaggttc	ggggggtgga	atttgatgac	aaagataagc	1860
agttcttcaa	aaactttgcg	acaggtgctg	ttcctatagc	atggcaggaa	gaaattatag	1920
aaacgggact	gtttgaggaa	ctgaatgacc	ccaacagacc	tacgggttgt	gaggagggta	1980
attcatccaa	gtctggcgtg	tgtttgttat	tgtaaatgtc	tctctttacc	agacaggcag	2040
caggagtctc	ggctgacata	atcctcgaat	gttccacacg	tggaatctg	tggaatgagg	2100
gctaatacagt	taggaggggac	atcacaacca	caaaacaatt	caaaagacag	gcaagctcac	2160
tactagaaca	cattttattt	tctttttctt	tcttcataaa	gatgagtaaa	gtctcagttt	2220
tcactgaggg	cagggaaaaag	gaacactca				2249